



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Surgery for patients with 'lower grade' glioma: putting assumptions, beliefs and convictions into perspective

Weller, M

DOI: <https://doi.org/10.1093/annonc/mdx420>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-141079>

Journal Article

Accepted Version

Originally published at:

Weller, M (2018). Surgery for patients with 'lower grade' glioma: putting assumptions, beliefs and convictions into perspective. *Annals of Oncology*, 29(4):1077.

DOI: <https://doi.org/10.1093/annonc/mdx420>

Surgery for patients with “lower grade” glioma: putting assumptions, beliefs and convictions into perspective

Michael Weller

Department of Neurology, University and University of Hospital Zurich, Zurich,
Switzerland

For decades neuro-oncology meetings have benefited from controversial, sometimes heated and emotional debates on the role of surgery for what has traditionally been falsely designated “low grade” glioma; falsely since “low grade” glioma evokes false expectations for often young patients and their caregivers confronted with a disease with a median survival around 7-8 years.

Advocates of either radical surgery for almost everybody or surgery for almost nobody have repeatedly stated their concerns that randomized clinical trials in that situation would be unethical: what more evidence would we need to define a clinical trial as ethical if there is so little agreement among the “experts”?

Yet, in the absence of randomized data to address this question and in the absence even of efforts to do obtain such data, the Norwegian study, facilitated by a small population living in a big country with apparently little patient mobility at the time that the study was conducted, is one of the best estimates of the potential impact of early efforts at resection in patients with World Health Organization (WHO) grade II gliomas. In 2012, Jakola and colleagues reported their first comparison of two management strategies for patients with these tumors in two regions of Norway: one hospital (A) preferred a wait and scan strategy after biopsy whereas another hospital (B) favored early resections. In the initial report [1], median survival was 5.9 years for patients initially managed in hospital A whereas it was not reached in hospital B.

Estimates for 5-year survival were 60% versus 74%. Multivariable analyses determined the relative hazard ratio for death to be 1.8 when a patient was treated in hospital A as opposed to hospital B.

In the present update [2], long-term survival data are reported: median overall survival was 5.8 years in hospital A as opposed to 14.4 years in hospital B.

Importantly, the authors undertook a recommendable effort to provide a molecular marker-based stratification of their cohorts according to the new WHO classification and determined that the potential impact of efforts at early resection remained after adjustment for molecular phenotype: IDH mutant 1p/19q codeleted versus IDH mutant versus IDH wild-type.

Admittedly, the study has several limitations which include, amongst others, that with a median age of 44-45 years in both arms, at least half of the patients were in an “elder” age group where most centers would probably not advocate a wait-and-scan strategy in the first place, moreover, the molecular data suffer from the limitation that these data were not assessed centrally or according to the same methodology.

The extensive literature on the role of surgery in gliomas suffers from many assumptions, overinterpretations and misconceptions, and many publications are overtly driven by bias [3,4]: for instance, numerous publications claim *effect* or *impact* of surgery for gliomas although non-randomized retrospective cohort studies [5-7] can detect associations with outcome at best, but by definition can never demonstrate effect or impact (although the Norwegian data set comes relatively close to that).

Similar considerations apply to the discussion of surgery for WHO grade IV gliomas (glioblastomas). There is an almost bizarre discussion on estimating the extent of resection in precise percentages [8] while logics tell us that residual tumor volume – rather than extent of resection - should be in the focus [9], irrespective how big the

tumor, glioma of either WHO grade II, III or IV, was on preoperative scans. A 90% resection of a 100 mm³ tumor still leaves a lesion of double size relative to a 50% resection of a 10 mm³ tumor.

Given the plethora of publications that have not advanced the controversy in this area to a relevant extent, the Norwegian data likely provides (one of) the strongest pieces of evidence to support a role for early surgical interventions in patients with so called “low grade” gliomas. Importantly, it indicates a role of surgery across the three major molecular phenotypes of these tumors defined by the 2016 WHO classification. This does not contradict, but rather complements the recent research focus on whether and how molecular markers determine resectability of gliomas [10].

Where do we go from here? Will the present study [1,2] remain to be the best evidence to support a role for surgery in WHO grade glioma? While extent and also timing of surgery remain central topics in clinical research on these tumors, the WHO classification has started to dissect this group of tumors into smaller entities, and new questions arise, notably on targeted systemic treatments e.g. focusing on mutant isocitrate dehydrogenase either as a pharmacological or an immunotherapeutic target. In that emerging landscape of clinical trials, it is increasingly unlikely that the question of surgery will ever be answered in a much more conclusive way than here [1,2].

There was no funding for this work and the author has declared no conflict of interest.

References

1 Jakola AS, Myrnes KS, Kloster R et al. Comparison of a strategy favoring early surgical resection vs a strategy favoring watchful waiting in low-grade gliomas. JAMA 2012; 308: 1881-1888.

2 Jakola AS, Skjalsvik AJ, Myrnes KS et al. Surgical resection versus watchful waiting in low-grade gliomas. Ann Oncol 2017; XX: YYY-YYY.

3 Jiang B, Chaichana K, Veeravagu A, Chang SD, Black KL, Patil CG. Biopsy versus resection for the management of low-grade gliomas. Cochrane Database Syst Rev 2017; 27: 4:CD009319.

4 Weller M, van den Bent M, Tonn JC et al., for the European Association for Neuro-Oncology (EANO) Task Force on Gliomas. EANO guideline on the diagnosis and treatment of adult astrocytic and oligodendroglial gliomas. Lancet Oncol 2017 in press.

5 Smith JS, Chang EF, Lamborn KR et al. Role of extent of resection in the long-term outcome of low-grade hemispheric gliomas. J Clin Oncol 2008; 26: 1338-1345.

6 Capelle L, Fontaine D, Mandonnet E et al. Spontaneous and therapeutic prognostic factors in adult hemispheric World Health Organization Grade II gliomas: a series of 1097 cases. J Neurosurg 2013; 118: 1157–1168.

7 Eseonu CI, Eguia F, ReFaey K et al. Comparative volumetric analysis of the extent of resection of molecularly and histologically distinct low grade gliomas and its role on survival. J Neurooncol 2017 [Epub ahead of print]

8 Lacroix M, Abi-Said D, Fourney DR, et al. A multivariate analysis of 416 patients with glioblastoma multiforme: prognosis, extent of resection, and survival. J Neurosurg 2001; 95: 190-198.

9 Chaichana KL, Jusue-Torres I, Navarro-Ramirez R et al. Establishing percent resection and residual volume thresholds affecting survival and recurrence for patients with newly diagnosed intracranial glioblastoma. Neuro Oncol 2014; 16: 113-122.

10 Beiko J, Suki D, Hess KR et al. IDH1 mutant malignant astrocytomas are more amenable to surgical resection and have a survival benefit associated with maximal surgical resection. Neuro-Oncology 2014; 16: 81–91.